

TITLE PAGE

Management of a Hyperactive Gag Reflex for OAT: A Case Report

Andrew S. Ryser, CPT, DDS, ^a Kyung Kim, COL, DMD, ^b Steven Handel, LTC, DMD, ^c
Jennifer Sabol, COL, DDS, ^d Maddie E. Ryser, DNP, FNP-BC, ^e

^a Prosthodontic Resident, Department of Prosthodontics, Army Postgraduate Dental School,
Postgraduate Dental College, Uniformed Services University, Fort Eisenhower, GA, USA

^b Assistant Program Director, Department of Prosthodontics, Army Postgraduate Dental School,
Postgraduate Dental College, Uniformed Services University, Fort Eisenhower, GA, USA

^c Assistant Program Director, Department of Prosthodontics, Army Postgraduate Dental School,
Postgraduate Dental College, Uniformed Services University, Fort Eisenhower, GA, USA

^d Program Director, Department of Prosthodontics, Army Postgraduate Dental School,
Postgraduate Dental College, Uniformed Services University, Fort Eisenhower, GA, USA

^e Private Practice, GA

DISCLAIMER: The views expressed in this manuscript are those of the authors and do not necessarily reflect the official policy of the United States Government, the Department of Defense, the Defense Health Agency, or Uniformed Services University.

This research did not receive any specific grant from funding agencies in the public, commercial, or not for-profit sectors.

CASE REPORT

Management of a Hyperactive Gag Reflex for OAT: A Case Report

ABSTRACT

Background: A patient sought treatment at a prosthodontics practice for treatment of obstructive sleep apnea (OSA) with a mandibular advancement device (MAD).

Case Information: Due to a hyperactive gag reflex, a patient required implementation of a unique protocol to overcome gagging whenever the MAD was placed in the mouth.

Conclusion: The protocol described in this case report allowed the patient to overcome a hyperactive gag reflex and successfully use a MAD to treat OSA.

INTRODUCTION

Impression taking and other dental procedures such as wearing a maxillary complete denture can be stressful for some patients and dentists due to the presence of a gag reflex. Sivakumar and Prabhu defined the gag reflex as an involuntary reflex involving bilateral pharyngeal muscle contraction and elevation of the soft palate.¹ It is believed that the gag reflex arose as humans evolved to prevent us from swallowing foreign objects and choking.¹ Physical examination of reflexes such as the gag reflex and response to pain are often used to determine brain activity levels or if brain death has occurred.² The gag reflex is controlled by cranial nerves IX (glossopharyngeal nerve) and X (vagus nerve) and is derived from the third, fourth, and sixth pharyngeal arches during development.¹ In newborns, the gag reflex is activated whenever the brainstem deems food to be too large or solid for an infant to digest. At around a half year of age, the gag reflex becomes less pronounced, and the infant can start consuming more solid foods.³

The gag reflex can either be classified as somatogenic or psychogenic in nature. A somatogenic gag reflex occurs when there is direct physical contact with the base of tongue, the tonsillar pillar area, or the posterior pharyngeal wall with a physical object such as a piece of food, dental hand mirror, or oral appliance. A psychogenic gag reflex occurs from a mental trigger, often without a physical contact. A psychogenic factor could be the simple thought of going to a dental appointment inducing the gag reflex. ⁴

A direct gag reflex is defined as the contraction of the pharyngeal musculature ipsilateral to the side of the stimulus. A consensual gag reflex is when the musculature of the pharynx contracts on the contralateral side of the stimulus.¹ Gagging can also occur when the soft palate is stimulated by activating CN V (trigeminal nerve). According to Davies et al. in 1995, around 37% of people do not have a gag reflex.⁵ Davies and co-authors also concluded that there is an increase in the absence of a gag reflex in those that smoke or use tobacco.⁵ An absent gag reflex can be a cause for concern however as it might indicate cranial nerve damage or brain death.⁵ Neumann and McCarty in 2001 found that 10-15% of people have a hypersensitive gag reflex and those people will often gag when eating sticky or thick foods.⁶

There are many etiologic factors that stimulate the gag reflex to include:

- 1) Local factors: nasal obstruction, deviated septum, postnasal drip, sinusitis, congestion.
- 2) Systemic conditions: chronic gastritis, Paterson's dysplasia, stomach and pancreas cancer, uncontrolled diabetes and peptic ulcers.
- 3) Social factors: smoking, coughing, chronic alcoholism.
- 4) Psychological factors: eating disorders, stress, neuroticism.

- 5) Iatrogenic factors: medical/ dental instruments, local anesthesia, small tongue space.
- 6) Prosthetic Factors: poor retention of prostheses, overextended or underextended dentures, inadequate posterior palatal seal.⁷⁻⁹

The Dickinson and Fiske gagging severity index can be used to classify the gag reflex severity and is as follows:¹⁰

Grade 1: Very mild: patient is able to control gagging

Grade 2: Mild: gagging is able to be controlled by patient/ dentist using simple measures

Grade 3: Moderate: some treatment is not accepted by patient without gagging

Grade 4: Severe: some treatment is not possible

Grade 5: Very severe: any procedure in the oral cavity is impossible and effects patient behavior.

Providers can test for a gag reflex by using a tongue blade or cotton tip applicator. Using one of the two instruments, the provider stimulates each side of the posterior pharynx waiting each time for gagging.¹ In patients with severe gagging placing dental instruments such as impression trays, a gloved finger, or even intraoral mirrors could be enough to stimulate the gag reflex. Management techniques to control gagging include behavior modification, desensitization, conditioning prostheses, pharmacological therapy, acupuncture/ acupressure, modification of prostheses, hypnosis, and holistic remedies.¹¹ This article discusses a new and novel technique used on a patient with a Dickinson and Fiske Grade 4 gag reflex to successfully treat the patient's obstructive sleep apnea (OSA) with a mandibular advancement device.

NARRATIVE

A 60-year-old patient presented to a prosthodontics practice for treatment of OSA and bruxism. The patient was 64 inches tall and weighed 155 pounds with a BMI of 26.8. The patient's medical history was significant for hypertension, hyperlipidemia, Parkinson's, impaired fasting glucose, melanoma, allergic rhinitis, somatoform disorder, and obstructive sleep apnea. A full night polysomnogram (PSG) at Fort Eisenhower sleep clinic was completed, and the PSG results revealed an AHI of 12.9 with moderate to loud snoring. The total sleep time of the PSG was 362 minutes, with 11% of total sleep in stage N1, 44.1% in Stage N2, 32.6% in Stage N3, and 12.3% in REM. Sleep latency was 9.5 minutes and REM latency was 119 minutes. The patient spent 48.5 minutes sleeping in the supine position. Baseline oxygen saturation was 94.9% and the lowest recorded oxygen saturation was 87.9%. Increased periodic limb movements were observed.

The patient completed an in-home auto titration study that demonstrated effective treatment of OSA, but it also demonstrated inappropriate compliance using CPAP. The patient followed up with the sleep clinic stating she did not want to use a CPAP and wanted to try oral appliance therapy. The patient's intraoral exam revealed past dental restorations, attrition of the dentition, mild crowding, barely visible tonsils, a normal uvula size, and a class 3 Mallampati score. The patient's Epworth sleepiness scale score was fifteen and she complained of snoring, daytime drowsiness, bruxism, drooling while sleeping, and restlessness.

The patient had a hyperactive gag reflex which presented anytime the intraoral mirror would touch the tongue or go past the anterior dentition. Intra-oral scans with a Primescan (Dentsply Sirona, USA) were obtained by placing salt on the tongue as described by Friedman.¹⁴ A protrusive bite registration using a George Gauge with 3 mm thick bite fork (Great Lakes Sleep Technologies, USA) at 60% maximum protrusion was made. The scans and bite

registration were sent to the lab and an EVO (Prosomnus Sleep Technologies, USA) appliance was fabricated. During the delivery appointment, the patient was unable to tolerate the maxillary and mandibular EVO appliance without gagging. A desensitization protocol was developed and implemented to aid the patient in being able to utilize the MAD to treat her OSA. The protocol was as follows:

- Day 1-Day 7: The patient would wear the maxillary MAD for an increasing amount of time throughout the day as much as the patient could tolerate.
- Day 8-Day 14: The patient would use both maxillary and mandibular trays simultaneously for an increasing amount of time throughout the day.
- Day 15: The patient would use the MAD at bedtime and go to sleep while using it for as long as tolerable.
- Day 16: The patient stated she was able to lay in bed and go to sleep with the MAD but after a brief period she would start gagging intermittently. The patient also stated she could sit up without any gagging occurring. The protocol was modified, and the patient was instructed to go to sleep with three pillows propping her head/upper body up.
- Day 18: The patient was instructed to take one pillow away
- Day 20: The patient was instructed to take the second pillow away. The patient was able to successfully lay in bed using the MAD without gagging and without any positional modification to elevate head/upper body.

The patient returned to the clinic for re-evaluation about 21 days after the desensitization period ended. The patient stated she was able to use the MAD on a regular basis and was able to sleep throughout the night with it in. She reported getting better sleep and waking up in the morning feeling more energetic and that she was feeling less tired during the day. The patient

was referred to her PCM for a follow up sleep study, however the MAD efficacy PSG has not yet been completed due to urgent medical conditions the patient has experienced. The patient and PCM are planning a PSG once the patient's medical condition has stabilized.

DISCUSSION

According to Bassi et al., gagging can be disruptive to dental treatment and can prevent the wearing of a prosthesis such as a complete denture or a mandibular advancement device.⁷ Gagging can be trying for the patient and make dentistry difficult or sometimes even impossible to perform. As previously stated, many different techniques have been developed to prevent gagging to include medications, sedatives, holistic remedies such as herbs, acupressure and acupuncture, sedatives, alteration of prosthetic devices, and general or local anesthesia.¹¹

Often simple behavior modification and distraction such as providing reassurance and having the patient attempt to relax with music, television, or talking can stop the gag reflex.⁹ Pharmacological options to reduce the gag reflex include local anesthetic, conscious sedation, and general anesthesia.^{12,13} Altering the size and shape of a prosthesis, such as reduction of the palate on a maxillary complete denture could be enough to prevent gagging.¹⁴ Other prosthodontic management techniques for gagging include: reducing thickness and extensions of prostheses, choosing impression material with a high viscosity, giving patients training prostheses prior to final prosthesis delivery, and making denture surfaces with a matte finish instead of smooth and glossy.¹⁵ Friedman discussed placing salt on the palatal region of a prosthesis or on the tongue to reduce gagging; a technique often used in dentistry to reduce gagging when taking radiographs or making impressions.¹⁶ In 2022, Mehdizadeh et al. performed a systematic review that provided an update on the management of the gag reflex and stated that

nitrous oxide, low level lasers, and distraction techniques were effective in reducing the occurrence of the gag reflex.¹⁷

A common technique used in dentistry to control the gag reflex is desensitization. Desensitization consists of slowly exposing the patient to a stimulus that invokes the gag reflex. Frequency, duration, and intensity of the stimulus is gradually increased with the goal of habituating the procedure that is to be performed or the use of a prosthesis such as a complete denture or other removable prosthesis.⁹ Colvenkar et al. described a step by step process to manage gagging in an edentulous patient which included having the patient brush his palate as far posterior as possible with a toothbrush and placing 4 pieces of chocolates in his mouth one by one until all four pieces would go in without gagging.¹⁵ In 1973 Singer published an article discussing desensitization for treating patients with a severe gag reflex. The Singer's marble technique called for placing marbles in the patient's mouth to desensitize the patient in preparation for use of complete dentures, demonstrating an odd but novel way to treat patient successfully.¹⁸ In dentistry providers often must come up with creative ways to desensitize/distract, as Colvenkar and Singer did, to perform procedures that might stimulate the gag reflex, calm fears, or decrease anxiety.

CONCLUSION

This case report discusses a novel technique used to help a patient acclimate to using a mandibular advancement device without activation of the gag reflex. The described protocol works with the patient's tolerances for wearing the appliance in small increments, allowing the patient to have complete control of the environment. The result allowed the patient to be able to utilize the MAD without the occurrence of gagging. This new and novel technique was

successful for this patient, has the potential to help other patients overcome a hyperactive gag reflex, and can be personalized for each individual patient.

REFERENCES

1. Sivakumar S, Prabhu A. Physiology, Gag Reflex [Internet]. PubMed. Treasure Island (FL): Stat Pearls Publishing; 2020.
2. Park MJ, Byun JS, Jung JK, Choi JK. The correlation of gagging threshold with intra-oral tactile and psychometric profiles in healthy subjects: A pilot study. *J Oral Rehabil.* 2020;47:591-598.
3. Stevenson RD, Allaire JH. The development of normal feeding and swallowing. *Pediatr Clin North Am.* 1991;38:1439-53.
4. Saunders RM, Cameron J. Psychogenic gagging: identification and treatment recommendations. *Compend Contin Educ Dent.* 1997;18:430-3, 436, 438.
5. Davies AE, Kidd D, Stone SP, MacMahon J. Pharyngeal sensation and gag reflex in healthy subjects. *Lancet.* 1995;345:487-8.
6. Neumann JK, McCarty GA. Behavioral approaches to reduce hypersensitive gag response. *J Prosthet Dent.* 2001;85:305
7. Bassi GS. The Etiology and Management of Gagging. A Review of the Literature. *J Prosthet Dent* 2004;9:459.
8. Lavinia Ardelean, et al. Gag Reflex in Dental Practice Etiological Aspects. *Tmj* 2003, Vol. 53, No. 3-4
9. Shrivastav VK, Mahajan S. Effective management of gag reflex in clinical practice. *Indian J Stomatol* 2012;3:27-31.

10. Pisulkar SK, Agrawal R, Godbole SR, Jadhav V. Addressing the gag reflex: A literature review. *International Journal of Recent Surgical and Medical Sciences*. 2018;04:002–4.
11. Eachempati P, Kumbargere Nagraj S, Kiran Kumar Krishanappa S, George RP, Soe HHK, Karanth L. Management of gag reflex for patients undergoing dental treatment. *Cochrane Database of Systematic Reviews*. 2019
12. Braham RB, Braham RL. The management of the chronic oropharynx and palate. *ASDC J Dent Child* 1977;44:111-116.
13. Yagiela JA. Making patients safe and comfortable for a lifetime of dentistry: frontiers in office-based sedation. *J Dent Educ* 2001;65:1348-1356.
14. Borkin DW. Impression technique for patients that gag. *J Prosthet Dent* 1959;9:386-387.
15. Colvenkar S, Reddy V, Thotapalli S, Deepa Rani K, Bharadwaj S. A Simple Step-by-Step Technique for the Management of Gagging in Edentulous Patient. *Cureus*. 2022;14:e24423.
16. Friedman MH, Weintraub MI. Temporary elimination of gag reflex for dental procedures. *J Prosthet Dent* 1995;73:319.
17. Mehdizadeh, M., Mohammadbeigi, A., Sharifinejad, A. An Overview about New Methods in Management of Gag Reflex during Dental Treatment: A Systematic Review. *Journal of Dentistry*, 2022
18. Singer IL. The marble technique: a method for treating the “hopeless gagger” for complete dentures. *J Prosthet Dent* 1973;29:146-150.

